

## CYC Vaccination Policy – DRAFT

As CYC is a intentional community of Christian fellowship that resides in close physical contact during the annual conference, health and safety concerns are of major importance. In view of this, it is the policy of CYCOP to require all of its participants (students, staff, faculty, and attending family members of staff and faculty) to adhere to the immunization regulations for summer camps set forth in section 105 of the Commonwealth of Massachusetts Regulations 430.152, unless the participant has a medical exemption certified by a physician. As of 2022, those requirements are:

### Participants in Grades K-6 (staff/faculty children)

DTaP	<b>5 doses;</b> 4 doses are acceptable if the 4 <sup>th</sup> dose is given on or after the 4 <sup>th</sup> birthday. DT is only acceptable with a letter stating a medical contraindication to DTaP.
Polio	<b>4 doses;</b> 4 <sup>th</sup> dose must be given on or after the 4 <sup>th</sup> birthday and ≥6 months after the previous dose, or a 5 <sup>th</sup> dose is required. 3 doses are acceptable if the 3 <sup>rd</sup> dose is given on or after the 4 <sup>th</sup> birthday and ≥6 months after the previous dose.
Hepatitis B	<b>3 doses;</b> laboratory evidence of immunity acceptable
MMR	<b>2 doses;</b> first dose must be given on or after the 1 <sup>st</sup> birthday and the 2 <sup>nd</sup> dose must be given ≥28 days after dose 1; laboratory evidence of immunity acceptable
Varicella	<b>2 doses;</b> first dose must be given on or after the 1 <sup>st</sup> birthday and 2 <sup>nd</sup> dose must be given ≥28 days after dose 1; a reliable history of chickenpox* or laboratory evidence of immunity acceptable

### Participants in Grades 7-12

Tdap	<b>1 dose;</b> and history of DTaP primary series or age appropriate catch-up vaccination. Tdap given at ≥7 years may be counted, but a dose at age 11-12 is recommended if Tdap was given earlier as part of a catch-up schedule. Td or Tdap should be given if it has been ≥10 years since Tdap.
Polio	<b>4 doses;</b> 4 <sup>th</sup> dose must be given on or after the 4 <sup>th</sup> birthday and ≥6 months after the previous dose, or a 5 <sup>th</sup> dose is required. 3 doses are acceptable if the 3 <sup>rd</sup> dose is given on or after the 4 <sup>th</sup> birthday and ≥6 months after the previous dose.
Hepatitis B	<b>3 doses;</b> laboratory evidence of immunity acceptable. 2 doses of Heplisav-B given on or after 18 years of age are acceptable.
MMR	<b>2 doses;</b> first dose must be given on or after the 1 <sup>st</sup> birthday and the 2 <sup>nd</sup> dose must be given ≥28 days after first dose; laboratory evidence of immunity acceptable
Varicella	<b>2 doses;</b> first dose must be given on or after the 1 <sup>st</sup> birthday and 2 <sup>nd</sup> dose must be given ≥28 days after first dose; a reliable history of chickenpox* or laboratory evidence of immunity acceptable
MenACWY (formerly MCV4)	<b>Grades 7: 1 dose;</b> MenACWY required. <b>Grades 11: 2 doses;</b> 2 <sup>nd</sup> dose MenACWY must be given on or after the 16 <sup>th</sup> birthday and ≥ 8 weeks after the previous dose. 1 dose is acceptable if it was given on or after the 16 <sup>th</sup> birthday. Meningococcal B vaccine is not required and does not meet this requirement.

### Participants 18 years of age and older\*

MMR	<b>2 doses,</b> anyone born in or after 1957. 1 dose, anyone born before 1957 outside the U.S. Anyone born in the U.S. before 1957 is considered immune. Laboratory evidence of immunity to measles, mumps and rubella is acceptable
Varicella	<b>2 doses,</b> anyone born in or after 1980 in the U.S., and anyone born outside the U.S. Anyone born before 1980 in the U.S. is considered immune. A reliable history of chickenpox** or laboratory evidence of immunity is acceptable
Tdap	<b>1 dose;</b> and history of DTaP primary series or age appropriate catch-up vaccination. Tdap given at ≥7 years

	may be counted, but a dose at age 11-12 is recommended if Tdap was given earlier as part of a catch- up schedule; Td or Tdap should be given if it has been $\geq 10$ years since Tdap
Hepatitis B	<b>3 doses (or 2 doses of Hepelisav-B) for staff whose responsibilities include first aid;</b> laboratory evidence of immunity is acceptable

\*Adult individuals who do not have access to their full immunization record may sign a form attesting to their having received the required immunizations.

\*\*A reliable history of chickenpox includes a diagnosis of chickenpox, or interpretation of parent/guardian description of chickenpox, by a physician, nurse practitioner, physician assistant or designee.